

## Statin Therapy Measures Tip Sheet

### Statin Therapy for Pts with Diabetes (SPD)

**Measure Description:** The % of patients 40-75yrs with a diagnosis of Diabetes during the measurement year who do not have Atherosclerotic Cardiovascular Disease (ASCVD).

**Measure Goal:** Dispensed at least one statin therapy of any intensity during the measurement year.

**Inclusion Criteria:** Patient medical claim and or pharmacy data. Any LDL result is inclusive.

### Statin Therapy for Pts with Cardiovascular Disease (SPC)

**Measure Description:** The % of males 21-75yrs and females 40-75yrs with a diagnosis Atherosclerotic Cardiovascular Disease (ASCVD).

**Measure Goal:** Dispensed at least one statin therapy of moderate or high intensity during the measurement year.

**Inclusion Criteria:** Patient with MI, CABG, PCI, or diagnosis of Ischemic Vascular Disease (IVD). Any LDL result is inclusive.

### Statin Use for Pts with Diabetes (SUPD)

**Measure Description:** The % of Medicare patients 40-75yrs with a diagnosis of Diabetes.

**Measure Goal:** Dispensed at least one statin therapy med of any intensity during the measurement year.

**Inclusion Criteria:** Patient medical pharmacy claim data only (at least 2 diabetes medication fills)

**Exclusions:** ESRD or Hospice.

Moderate Intensity Statin		High Intensity Statin	
Name of Drug	Dosage	Name of Drug	Dosage
Atorvastatin (Lipitor)	10-20 mg	Atorvastatin	40-80 mg
Amlodipine-atorvastatin (Caduet)	10-20 mg	Amlodipine-atorvastatin	40-80 mg
Ezetimibe-simvastatin (Vytorin)	20-40 mg	Ezetimibe-simvastatin	80 mg
Fluvastatin (Lescol)	40-80 mg	Rosuvastatin	20-40 mg
Lovastatin (Mevacor)	40 mg	Simvastatin	80mg
Pitavastatin (Livalo)	2-4 mg		
Pravastatin (Pravachol)	40-80 mg		
Rosuvastatin (Crestor)	5-10 mg		
Simvastatin (Zocor)	20-40 mg		

### Codes to submit for Statin Intolerance (Use for SPD & SPC only)

**Exclusion Criteria:** Cirrhosis, Myalgia, Myositis, Myopathy, Rhabdomyolysis, ESRD, Pregnant, ≥66 living in institutional or SNF setting, ≥81 with frailty claim, dispensed dementia medication and hospice.

Cirrhosis	Muscular Pain & Disease	End Stage Renal Disease
<b>K70.30</b> Alcoholic cirrhosis of liver w/o ascites	<b>G72.0</b> Drug induced myopathy	<b>N18.5</b> Chronic Kidney Disease, stage 5
<b>K70.31</b> Alcoholic cirrhosis of liver w/ascites	<b>G72.2</b> Myopathy due to other toxic agents	<b>N18.6</b> End Stage Renal Disease
<b>K71.7</b> Toxic liver w/fibrosis & cirrhosis of liver	<b>G72.9</b> Myopathy, unspecified	<b>Z99.2</b> Dependence on Renal Dialysis
<b>K74.3</b> Primary biliary cirrhosis	<b>M60.80</b> Other myositis, unspecified site	
<b>K74.4</b> Secondary biliary cirrhosis	<b>M60.89</b> Other myositis, multiple sites	
<b>K74.5</b> Biliary cirrhosis, unspecified	<b>M60.9</b> Myositis, unspecified	
<b>K74.60</b> Unspecified cirrhosis of liver	<b>M62.82</b> Rhabdomyolysis	
<b>K74.69</b> Other cirrhosis of liver	<b>M79.1</b> Myalgia	
<b>P78.81</b> Congenital cirrhosis (of liver)	<b>M79.10</b> Myalgia, unspecified site	
<b>K70.0</b> Alcoholic fatty liver	<b>M79.18</b> Myalgia, other site	
<b>K70.9</b> Alcoholic liver disease unspecified	<b>G72.89</b> Other specified myopathies	
	<b>T46.6X5A</b> Adverse effect of Antihyperlipidemic & antiarteriosclerotic drug initial encounter (SUPD only)	