

## Statin Therapy Measures Tip Sheet

### Statin Therapy for Pts with Diabetes (SPD)

**Measure Description:** The % of patients 40-75yrs with a diagnosis of Diabetes during the measurement year who do not have Atherosclerotic Cardiovascular Disease (ASCVD). PPO

**Measure Goal:** Dispensed at least one statin therapy of any intensity during the measurement year.

**Inclusion Criteria:** Patient medical claim and or pharmacy data. Any LDL result is inclusive.

### Statin Therapy for Pts with Cardiovascular Disease (SPC)

**Measure Description:** The % of males 21-75yrs and females 40-75yrs with a diagnosis Atherosclerotic Cardiovascular Disease (ASCVD). PPO /Medicare

**Measure Goal:** Dispensed at least one statin therapy of moderate or high intensity during the measurement year.

**Inclusion Criteria:** Patient with MI, CABG, PCI, or diagnosis of Ischemic Vascular Disease (IVD). Any LDL result is inclusive.

### Statin Use for Pts with Diabetes (SUPD)

**Measure Description:** The % of Medicare patients 40-75yrs with a diagnosis of Diabetes. BCNA/MA PPO

**Measure Goal:** Dispensed at least one statin therapy med of any intensity during the measurement year.

**Inclusion Criteria:** Patient medical pharmacy claim data only (at least 2 diabetes medication fills)

#### Moderate Intensity Statin

Name of Drug	Dosage
Atorvastatin (Lipitor)	10-20 mg
Amlodipine-atorvastatin (Caduet)	10-20 mg
Ezetimibe-simvastatin (Vytorin)	20-40 mg
Fluvastatin (Lescol)	40-80 mg
Lovastatin (Mevacor)	40 mg
Pitavastatin (Livalo)	2-4 mg
Pravastatin (Pravachol)	40-80 mg
Rosuvastatin (Crestor)	5-10 mg
Simvastatin (Zocor)	20-40 mg

#### High Intensity Statin

Name of Drug	Dosage
Atorvastatin	40-80 mg
Amlodipine-atorvastatin	40-80 mg
Ezetimibe-simvastatin	80 mg
Rosuvastatin	20-40 mg
Simvastatin	80mg

Medications Excluded: Zetia, Repatha and Praluent

### Codes to submit for Statin Intolerance (Use for SPD & SPC only)

**Exclusion Criteria:** Cirrhosis, Myalgia, Myositis, Myopathy, Rhabdomyolysis, ESRD, Pregnant, ≥66 living in institutional or SNF setting, ≥81 with frailty claim, dispensed dementia medication and hospice.

#### Cirrhosis

**K70.30** Alcoholic cirrhosis of liver w/o ascites  
**K70.31** Alcoholic cirrhosis of liver w/ascites  
**K71.7** Toxic liver w/fibrosis & cirrhosis of liver  
**K74.3** Primary biliary cirrhosis  
**K74.4** Secondary biliary cirrhosis  
**K74.5** Biliary cirrhosis, unspecified  
**K74.60** Unspecified cirrhosis of liver  
**K74.69** Other cirrhosis of liver  
**P78.81** Congenital cirrhosis (of liver)  
**K70.0** Alcoholic fatty liver  
**K70.9** Alcoholic liver disease unspecified

#### Muscular Pain & Disease

**G72.0** Drug induced myopathy  
**G72.2** Myopathy due to other toxic agents  
**G72.9** Myopathy, unspecified  
**M60.80** Other myositis, unspecified site  
**M60.89** Other myositis, multiple sites  
**M60.9** Myositis, unspecified  
**M62.82** Rhabdomyolysis  
**M79.1** Myalgia  
**M79.10** Myalgia, unspecified site  
**M79.18** Myalgia, other site  
**G72.89** Other specified myopathies

#### End Stage Renal Disease

**N18.5** Chronic Kidney Disease, stage 5  
**N18.6** End Stage Renal Disease  
**Z99.2** Dependence on Renal Dialysis